

Characterisation and treatment of patients with menstrual migraine in real-world clinical practice

Jessica Ailani¹, Gisela Terwindt², Boryana Galabova³, Jessica Cirillo⁴, Aaron Jenkins³, Lucy Abraham³, Karin Hygge Blakeman⁵, James Jackson⁶, Sophie Barlow⁶, William Whitton⁶, Lucy Hancock⁶

¹ MedStar Georgetown University Hospital, Washington DC USA; ² Department of Neurology, Leiden University Medical Center, Leiden, The Netherlands; ³ Pfizer, Ltd., Tadworth, Surrey; ⁴ Pfizer, Inc., New York, NY USA; ⁵ Pfizer AB, Stockholm, Sweden; ⁶ Adelphi Real World, Bollington, United Kingdom.

INTRODUCTION

- Menstrual migraine (MM) occurs in 2 out of 3 female migraine patients¹.
- Patients with MM experience more frequent and severe attacks than patients without MM².
- Despite this, MM is underdiagnosed and patients with MM are often treated similarly to migraine patients without MM.
- The objective of this study was to assess the demographics, clinical characteristics, treatment patterns and satisfaction of migraine patients with and without MM.

METHODS

- Data Source** – Data were drawn from the 2022/23 Adelphi Migraine Disease Specific Programme™, a real-world cross-sectional survey with retrospective data collection, conducted in France, Germany, Italy, Spain, the United Kingdom, and the United States. Physicians completed patient record forms for their next 8–10 consecutively consulting patients with migraine, who were invited to voluntarily complete a self-reported questionnaire.
- Study Design** – Participants were recruited into the Migraine DSP between May 2022 and January 2024. The survey was conducted according to relevant guidelines and legislation, and the methodology has been published and validated³⁻⁶.
- Analysis** – This analysis utilized data from patients who were female, aged ≤55 years old and had a physician confirmed diagnosis of migraine. Analysis was conducted on patient demographics, migraine diagnosis, including the estimated presence or absence of MM, prescribed acute and preventive treatment, and acute and preventive treatment satisfaction.
- Patients with reported MM were compared to patients without MM using: Mann-Whitney U tests for ordered categorical data, t-test for continuous outcomes, and Fisher exact tests for nominal categorical data. All data analyses were run using Stata 18⁷.

RESULTS


Patient demographics and clinical characteristics

- Primary care practitioners (n=264) and neurologists (n=340) provided data on 3,049 female patients aged ≤55 years old, of which 32% (n=986) were reported as having MM.
- Of these 3,049 female patients, 912 (30%) completed a corresponding voluntary self-reported questionnaire.
- Patients with reported MM were younger than patients without MM and experienced more severe migraine attacks as reported by physicians (all p<0.05, **Table 1** and **Figure 1**).

Table 1. Patient demographics and clinical characteristics			
	Patients with MM	Patients without MM	p-value
Patient age, n			
Mean (SD)	32.9 (9.0)	36.5 (10.1)	<0.0001 (TT)
BMI (≤60), n			
Mean (SD)	23.5 (3.9)	24.4 (4.6)	<0.0001 (TT)
Employment status, n (%)			
Working full time	570 (58.8)	1183 (59.7)	
Working part time	91 (9.4)	240 (12.1)	
On long term sick leave	8 (0.8)	25 (1.3)	
Homemaker	67 (6.9)	217 (11.0)	
Student	201 (20.7)	194 (9.8)	
Not working due to retirement	1 (0.1)	12 (0.6)	
Unemployed	31 (3.2)	111 (5.6)	

Abbreviations: TT; T test, MM, menstrual migraine; SD, standard deviation; BMI, body mass index
Base sizes vary due to availability of data in patient medical records. BMI Scores ≤60 were excluded.


Figure 1. Physician and patient-reported attack severity



CONFLICTS OF INTEREST

JA: Consulting (Honoraria): AbbVie, Aeon (2023), Dr. Reddy, Eli-Lilly, GlaxoSmithKline (2023), Lundbeck, Linpharma, Ipsen (2024), Merz, Miravio (2023), Pfizer, NeuroRelief, Gore, Satsuma (2024), Scilex, Theranica, Tonix. Clinical Trials (Grant to institution): Parema, Ipsen. Editorial Boards/Steering Committee: Current Pain and Headache Reports (2022), Medscape, SELF magazine (medical editor) GT: consultancy or industry support from AbbVie, Lilly, Lundbeck, Novartis, Pfizer, Teva, and Interactive Studios BV, and independent support from the European Community, Dutch Heart and Brain Foundations, Dutch Research Council, Biogen, and the Clayco Foundation. BG, JC, AJ, LA, KHB: employed by and holds stock/options in Pfizer. JJ, SB, WW, LH: employees of Adelphi Real World, Bollington UK.

Figure 2. Physician-reported current prescribed acute treatment



Currently prescribed treatments

- Patients without MM were more likely than patients with MM to be receiving no prescribed acute drug currently (p<0.05) (**Figure 2**).
- There was no differences between both groups for most acute treatments prescribed, with the exception of NSAIDs (p<0.05) (**Figure 2**).
- Patients with MM were more likely than patients without MM to be receiving no prescribed preventive drug currently (p<0.05) (**Figure 3**).
- Beta blockers and anti-CGRP mAbs were more likely to be prescribed for patients without MM (p<0.05) (**Figure 3**).

Treatment satisfaction

- Patients and physicians both reported a high level of satisfaction with treatment.
- Physicians reported a lower proportion of MM patients were extremely satisfied with their acute treatment compared to non-MM patients (**Figure 4**).
- Both physicians and patients reported a lower degree of satisfaction with preventive treatment for patients with MM. (**Figure 5**).

Figure 3. Physician-reported current prescribed preventive treatment




Figure 4. Physician and patient-reported satisfaction with acute treatment





Figure 5. Physician and patient-reported satisfaction with preventive treatment



Limitations

- MM was estimated by physicians and patients but not objectively assessed by a validated headache diary.
- While the study design required physicians to recruit consecutive patients and complete the physician survey on the day of the visit to mitigate against selection and recall bias. However, some selection bias remains as more frequently consulting patients and those with more severe disease activity were more likely to be captured.
- Disease severity data was for overall migraine severity, and not specifically within the perimenstrual window.

CONCLUSIONS

- While patients with and without menstrual migraine were prescribed similar treatment regimens, physicians estimated migraine attack severity to be worse and satisfaction with acute treatment lower in patients with MM.
- Education of physicians and their female patients is recommended to ensure optimal treatment for patients with MM and to improve satisfaction with therapy received.

REFERENCES

1: Burch *et al.* Headache: The Journal of Head and Face Pain, 2015;55(1):21-34. 2: Verhagen *et al.* Headache. 2023;63(3):333-41. 3: Anderson *et al.* Curr Med Res Opin. 2008;24(11):3063-3072. 4: Babineaux *et al.* BMJ Open. 2016;6(8):e010352. 5: Higgins *et al.* Diabetes Metab Syndr Obes. 2016;9:371-380. 6: Anderson P *et al.* Curr Med Res Opin. 2023;39(12):1707-1715. 7: StataCorp. 2023. Stata 18 Base Reference Manual. College Station, TX: Stata Press.

ACKNOWLEDGEMENTS

Data collection was undertaken by Adelphi Real World (Bollington, UK) as part of an independent survey, entitled the Adelphi Migraine Disease Specific Programme (DSP)™. All data are the intellectual property of Adelphi Real World. Pfizer Inc. subscribed to this survey and did not influence the original survey through either contribution to the design of questionnaires or data collection. Medical writing and editorial support was provided by Gary Sidgwick, PhD (Adelphi Real World, Bollington, UK), on behalf of Adelphi Real World and under the guidance of authors.



Electronic Poster: Please scan this Quick Response (QR) code with your smartphone app to view an electronic version of this poster. If you do not have access to a smartphone, please access the poster via the following link: <https://scientificpubs.congressposter.com/p/129v71227naeokfi>
Copies of this poster obtained through the QR code are for personal use only and may not be reproduced without permission from the authors of this poster.

Presented at the American Headache Society (AHS)
66th Annual Scientific Meeting,
June 13–15, 2024, San Diego, California, USA