

Long-term Efficacy and Safety of Lorlatinib vs. Alectinib in Patients With and Without Brain/Central Nervous System Metastases: Matching-Adjusted Indirect Comparisons

Objectives

- This study aimed to compare the long-term efficacy and safety of lorlatinib versus alectinib in previously untreated patients with anaplastic lymphoma kinase-positive (ALK+) metastatic non-small-cell lung cancer (mNSCLC) using matching-adjusted indirect comparisons (MAICs) based on the latest 5-year data from the CROWN trial (lorlatinib). Comparisons were conducted across three populations, including the intention-to-treat (ITT) population and patients with and without brain/central nervous system (CNS) metastases at baseline.

Conclusions

- This MAIC update provides long-term comparative data, further substantiating lorlatinib's effectiveness as a first-line treatment for ALK+ mNSCLC.
- Lorlatinib demonstrated superior long-term progression-free survival (PFS) compared with alectinib in the overall ITT population. Lorlatinib had a similar safety profile to alectinib in terms of adverse events (AEs) leading to treatment changes, although it was associated with a higher incidence of grade 3+ AEs.
- Lorlatinib significantly improved PFS in patients without baseline brain/CNS metastases over time and, despite the small sample sizes, demonstrated a numerical benefit over alectinib in patients with baseline brain/CNS metastases.



Click or scan this quick response (QR) to download this poster along with associated material.

References: 1. Ettinger DS, et al. / *Natl Comm Canc Netw*. 2022;20(5):497-530. 2. Peters S, et al. *N Engl J Med*. 2017;377(9):829-38. 3. Zhou C, et al. *Lancet Respir Med*. 2019;7(10):437-46. 4. Zhou C, et al. *Ann Oncol*. 2022;33:5158. 5. Mok T, et al. *Ann Oncol*. 2020;31(10):165-64. 6. Ahn MJ, et al. *Clin Lung Cancer*. 2022;23(8):720-30. 7. Camidge DR, et al. *J Thorac Oncol*. 2021;16(12):2001-106. 8. Garcia C, et al. *Clin Lung Cancer*. 2024;25(7):634-42. 9. Solomon BJ, et al. *Clin Oncol*. 2024;42(29):3400-9. 10. Shaw AT, et al. *N Engl J Med*. 2020;383(21):2018-29.

Acknowledgments: This study was sponsored by Pfizer. Evidera was a paid consultant for Pfizer in connection with the study and the development of this manuscript. Medical writing support was provided by Dr. Ruth Sharf (Evidera) and was funded by Pfizer, Inc. Disclosure: Devin Abraham, Anna Polli, Haitao Chu, and Priya Ramachandran are employees of Pfizer and have stock ownership in Pfizer. Todd Bauer receives consulting fees from and is an ad board member of Pfizer, Bayer, and Lilly. Julien Mazieres reports personal fees from Roche, Astra Zeneca, Pierre Fabre, Takeda, BMS, MSD, Pfizer, Jansing Hengshu, Blueprint, Daiichi, Novartis, Amgen and grants from Roche, Astra Zeneca, Pierre Fabre, BMS, Illumina. Christine Garcia receives consulting fees from Pfizer, Regeneron, Johnson and Johnson. Evidera received financial support from Pfizer in connection with the study and the development of this manuscript. Conor Chandler, Min Tan, and James Truscott are employees of Evidera, which was a paid consultant for Pfizer in connection with the study and the development of this manuscript.

Julien Mazieres¹, Todd Bauer², Devin Abraham³, Anna Polli³, Haitao Chu³, Priya Ramachandran³, Conor Chandler⁴, Min Tan⁴, James Truscott⁴, Christine A. Garcia⁵


¹Toulouse University Hospital, Institut Universitaire du Cancer, France; ²Department of Medical Oncology, Tennessee Oncology, Franklin, TN, USA; ³Pfizer Inc, New York, NY, USA; ⁴Evidera Inc, Bethesda, MD, USA; ⁵Department of Medicine, Weill Cornell Medicine and New York Presbyterian Hospital, New York, NY, USA

Background

- Over the past decade, the treatment landscape for ALK+ mNSCLC has evolved with the introduction of newer generations of ALK tyrosine kinase inhibitors (ALK TKIs).¹
- Second-generation (e.g., alectinib and brigatinib) and third-generation (e.g., lorlatinib) ALK TKIs have demonstrated greater efficacy as first-line treatments for ALK+ mNSCLC than crizotinib, a first-generation ALK TKI.²⁻⁷
- To date, no clinical trials have evaluated lorlatinib versus alectinib as first-line treatments for ALK+ mNSCLC, which necessitates alternative methods for comparative effectiveness research.
- An MAIC from an earlier 3-year data-cut of the CROWN trial showed that lorlatinib improved PFS compared with alectinib, although lorlatinib was associated with a higher rate of grade ≥3 adverse events (AEs).⁸
- The latest data-cut from October 31, 2023, allows for an updated long-term efficacy and safety analysis with extended follow-up data, including 5-year outcome data.

Results

- ITT Population (Figure 1)**
- Matching balanced differences in baseline characteristics across trials, reducing the effective sample size (ESS) by an acceptable ~15%. Lorlatinib demonstrated superior PFS compared with alectinib, reducing the risk of progression or death by 45% (HR: 0.55, 95% confidence interval [CI]: 0.34, 0.88).
 - Lorlatinib extended mean PFS by 8.5 months up to year 4 and by 11.2 months up to year 5.5, with an adjusted RMST approximately 1.5 to 1.6 times higher than alectinib.
 - PFS probabilities at years 1 to 5 were also significantly improved in terms of RDs with lorlatinib versus alectinib, ranging from 0.15 to 0.26 increases in annual PFS probabilities.
 - Lorlatinib had a higher rate of grade ≥3 AEs than alectinib. Rates of treatment discontinuation, dose interruption, and dose reduction were similar between lorlatinib and alectinib.




Methods

- An anchored MAIC was used to adjust for baseline differences in pre-selected effect modifiers, including the Eastern Cooperative Oncology Group Performance Score, presence of baseline brain/CNS metastases, and race (Asian vs. non-Asian).
- Published aggregate data for alectinib versus crizotinib were sourced from the ALEX⁹ trial, and patient-level data for lorlatinib versus crizotinib were sourced from the CROWN trial.^{9,10}
- Individual patient-level data from the CROWN trial were reweighted to match the average baseline characteristics of the patients in the ALEX trial, allowing for comparisons of outcomes across the two trials in similar populations.
- The MAIC analysis included both the ITT population and subpopulations with and without baseline brain/CNS metastases.

- Treatment effects for PFS were quantified using hazard ratios (HRs), as well as differences and ratios in annual probabilities and restricted mean survival time (RMST). AEs were compared based on rate ratios (RRs) and risk differences (RDs).
- RMST can be interpreted as the average PFS time during the period from baseline up to a specific timepoint of interest. In this analysis, RMST was measured up to year 4 and year 5.5 (end of follow-up in ALEX and CROWN).
- Observed Kaplan-Meier (KM) data were used where possible. However, parametric survival models were required to extrapolate PFS from years 2 to 4 for crizotinib (CROWN) in the subgroup with brain/CNS metastases. Due to crizotinib's limited efficacy in the brain, all patients either experienced PFS events or were censored before 24 months, necessitating the use of parametric models to estimate PFS beyond this period.

Population With Baseline Brain/CNS Metastases (Figure 2)

- Among patients with baseline brain/CNS metastases (ESS=64 of n=72), lorlatinib showed a benefit in PFS compared with alectinib, although this was not statistically significant because of the small sample size (HR: 0.47, 95% CI: 0.19, 1.20).
 - At year 1, lorlatinib significantly improved PFS (RD: 0.30, 95% CI: 0.03, 0.57), and numerical benefits were observed in subsequent years.
- Population Without Baseline Brain/CNS Metastases (Figure 2)**
- Among patients without baseline brain/CNS metastases (ESS=195 of n=204), lorlatinib demonstrated superior PFS compared with alectinib (HR: 0.51, 95% CI: 0.27, 0.94).
 - At year 1, lorlatinib was associated with a higher PFS compared with alectinib, although this result was not statistically significant. Over time, lorlatinib continued to show a consistent improvement in PFS, maintaining its advantage over alectinib and demonstrating a significant increase in PFS probabilities at years 2 to 4.



Abbreviations: CI = confidence interval; CNS = central nervous system; HR = hazard ratio; ITT = indirect treatment comparison; PFS = progression-free survival; PFS-NV = progression-free survival assessed by investigator; RD = risk difference; RR = rate ratio; W/O = without.